arXiv:1112.2811v2 [nucl-th] 28 May 2012

Real photons produced from photoproduction in pp collisions

FU Yong-Ping[1, 2] and LI Yun-De[1, *]

[1] *Department of Physics, Yunnan University, Kunming, 650091, China*
[2] *Department of Physics and Mathematics, Lincang Teachers College, Lincang, 677000, China*

We calculate the production of real photons originating from the photoproduction in relativistic pp collisions. The Weizsäcker-Williams approximation in the photoproduction is considered. Numerical results agree with the experimental data from Relativistic Heavy Ion Collider (RHIC) and Large Hadron Collider (LHC). We find that the modification of the photoproduction is more prominent in large transverse momentum region.

PACS: 25.20.Lj, 12.39.St, 12.38.Bx

Since photons do not participate in the strong interaction, the observation of prompt photons from the hadron collisions is a good method to test the perturbative Quantum Chromodynamics (pQCD). The application of pQCD for various hard scattering processes of the prompt photon production has been investigated [1–6]. It is necessary to identify different prompt photon sources in relativistic hadronic collisions. The hard scattering of partons is a well-known source of large transverse momentum (P_T) photons in high energy hadronic collisions. The prompt photons include direct photons and fragmentation photons [7–9]. The direct photons are those produced by the Compton scattering ($qg \to q\gamma$) and the annihilation of two partons ($q\bar{q} \to g\gamma$). The fragmentation photons are those produced by the bremsstrahlung emitted from final state partons ($ab \to (c \to x\gamma)d$). Previous works have studied the next-to-leading-order (NLO) contributions of the prompt photon production [10, 11], but the contributions of the photoproduction (higher-order) in the pp collisions are not researched clearly.

The photoproduction is an important concept in the ep deep inelastic scattering (DIS) at Hadron Electron Ring Accelerator (HERA) [12–14]. In the ep DIS, the electron interacts with the partons of the proton by exchanging a photon, a high energy photon emitted from the incident electron ($e \to e\gamma$) directly interacts with the proton by the interaction of $\gamma p \to$X (photoproduction). Besides, the Heisenberg's uncertainty principle allows the high energy photon for a short time to fluctuate into a quark-antiquark pair which then interacts with the partons of the proton (resolved photoproduction). In such interactions the photons emitted from the incident electron can be regarded as an extended object consisting of quarks and also gluons. The photons are the so-called resolved photons ($\gamma_{resolved}$).

In this work, we study the large P_T photons from the photoproduction mechanism by considering the Weizsäcker-Williams approximation in relativistic pp collisions. The charged partons of the incident proton can emit high energy photons ($q \to q\gamma$) and resolved photons ($q \to q\gamma_{resolved}$) in relativistic pp collisions [15–19], then the high energy photons or resolved photons interact with the partons of another proton by the subprocesses of $q\gamma \to q\gamma$, $q_\gamma\bar{q} \to g\gamma$, $\bar{q}_\gamma q \to g\gamma$, $q_\gamma g \to q\gamma$ and $g_\gamma q \to q\gamma$, here $q_\gamma(g_\gamma)$ denotes the parton of the resolved photon.

The prompt photons produced by the leading-order (LO) QCD Compton scattering, annihilation, and bremsstrahlung from final state partons have been discussed by previous works [1, 9, 10]. In this Letter we focus on the higher-order modification: the photoproduction. In the photoproduction, a parton q_a of an incident proton A emits a high energy photon ($q_a \to q_a\gamma$), then the photon interacts with a parton q_b of another incident proton B by the interaction of $\gamma p \to \gamma X$. The momentum transfer Q^2 in the deep inelastic collisions is large, so the perturbative factorization is applicable. The large P_T photons produced by photoproduction (pho.) satisfy the following invariant cross section [15, 16]

$$E\frac{d\sigma_{pho.}}{d^3P} = \frac{2}{\pi}\int_{x_{amin}}^{1} dx_a \int_{x_{bmin}}^{1} dx_b F_{q_a/A}(x_a, Q^2) \times F_{q_b/B}(x_b, Q^2) f_{\gamma/q_a}(z_a)\frac{x_a x_b z_a}{x_a x_b - x_a x_2} \times \frac{d\hat{\sigma}}{d\hat{t}}(x_a, x_b, z_a, P_T), \tag{1}$$

where x_a and x_b are the momentum fractions of partons, $F_{q_a/A}(x_a, Q^2)$ and $F_{q_b/B}(x_b, Q^2)$ are the parton distribution of the proton [20], we choose $Q^2 = 4P_T^2$ in the distribution [7]. $f_{\gamma/q_a}(z_a)$ is the photon spectrum from the quark q_a. The cross section of LO subprocesses $d\hat{\sigma}/d\hat{t}$ is QED Compton scattering $q_b\gamma \to q\gamma$. The minimum values of x_a and x_b in the integral are

$$x_{amin} = \frac{x_1}{1 - x_2}, \tag{2}$$

and

$$x_{bmin} = \frac{x_a x_2}{x_a - x_1}. \tag{3}$$

The momentum fraction z_a of the photon emitted from the quark is

$$z_a = \frac{x_b x_1}{x_a x_b - x_a x_2}, \tag{4}$$

where the variables are

$$x_1 = \frac{1}{2}x_T e^y, \tag{5}$$

---

[*] Corresponding author *E-mail address:* yndxlyd@163.com

and

$$x_2 = \frac{1}{2} x_T e^{-y}, \tag{6}$$

and

$$x_T = \frac{2P_T}{\sqrt{s_{NN}}}, \tag{7}$$

here $\sqrt{s_{NN}}$ is the total energy in the center-of-mass system, and y is the rapidity of the real photons.

The pQCD requests the QCD momentum transfer $Q^2 > 1$ GeV^2 [19]. In the photoproduction high energy photons are bremsstrahlung from the charged particles, this is a pure QED process. Since the QCD coupling parameter $\alpha_s(Q^2)$ does not depend on the QED momentum transfer q^2, the Weizsäcker-Williams approximation of the photoproduction is still valid for the charged partons of protons [18, 19]. Therefore the photon spectrum from the quark can be described by the Weizsäcker-Williams distribution

$$\begin{aligned} f_{\gamma/q_a}(z_a) = \int_{q^2_{min}}^{q^2_{max}} dq^2 \frac{dN_\gamma}{dzdq^2} = \\ \times \frac{e_q^2 \alpha P_{q\to\gamma}(z_a)}{2\pi} \left[\ln\left(\frac{(\sqrt{s_{NN}})^2}{4m_q^2} \right) + \ln\left(\frac{x_a x_b (1-z_a)}{z_a} \right) \right] \\ - \frac{e_q^2 \alpha}{\pi} \left(\frac{1-z_a}{z_a} - \frac{4m_q^2}{x_a x_b s_{NN}} \right), \end{aligned} \tag{8}$$

where e_q is the charge of the quark, α is the electromagnetic coupling parameter, $P_{q\to\gamma}(z_a) = \left(1 + (1-z_a)^2\right)/z_a$ is the split function. The maximum momentum transfer q^2_{max} is determined by the experimental acceptance [12]. According to [19] we choose q^2_{max} as $\hat{s}/4$. By considering the Weizsäcker-Williams approximation, the kinematic limit of minimum momentum transfer $q^2_{min} = (m_q z_a)^2/(1-z_a)$ is made such that the photon is close to being on its mass shell [13, 14]. Here m_q is the mass of the quark. The photoproduction contain higher-order QED coupling parameters, but the collision energies at RHIC ($\sqrt{s_{NN}} = 200$ GeV) and LHC ($\sqrt{s_{NN}} = 5500$ GeV) are large enough. Therefore the values of the term $\ln\left((\sqrt{s_{NN}})^2/4m_q^2\right)$ in Eq. (8) enhance the modification of the photoproduction [15].

The QED Compton process ($q_b\gamma \to q\gamma$) can be written as [7]

$$\frac{d\hat{\sigma}}{d\hat{t}}(q\gamma \to q\gamma) = \frac{\pi\alpha^2 e_q^4}{\hat{s}^2} 2 \left(-\frac{\hat{t}}{\hat{s}} - \frac{\hat{s}}{\hat{t}} \right), \tag{9}$$

where the Mandelstam variables are

$$\hat{s} = x_a x_b z_a s_{NN}, \tag{10}$$

and

$$\hat{t} = -x_a x_2 z_a s_{NN}. \tag{11}$$

The high energy photon emitted from the quark of the proton can fluctuate into a parton anti-parton pair in a short period of time $\triangle t$. Besides the (anti-)quarks may emit gluons in the time $\triangle t$. If, during such a fluctuation, one of the partons from the fluctuated photon interacts with partons of another incident proton. In such interaction the high energy photon is resolved [12–14]. In the resolved photoproduction, the parton q_a of the incident proton A emits a high energy resolved photon ($q_a \to q_a \gamma_{resolved}$), then the parton of the resolved photon ($\gamma_{resolved} \to q_{a'}\bar{q}_{a'}$ or $q_{a'}\bar{q}_{a'}g_{a'}$) interacts with the parton (q_b or g_b) of another incident proton B by the interaction of $\gamma_{resolved} p \to \gamma X$. The corresponding invariant cross section of large P_T photons produced by resolved photoproduction (res. pho.) can be written as [15, 16]

$$\begin{aligned} E\frac{d\sigma_{res.pho.}}{d^3P} = \frac{2}{\pi} \int_{x_{amin}}^{1} dx_a \int_{x_{bmin}}^{1} dx_b \int_{z_{a'min}}^{1} dz_{a'} \\ \times F_{q_a/A}(x_a, Q^2) F_{q_b/B}(x_b, Q^2) f_{\gamma/q_a}(z_a) \\ \times F_{q_{a'}/\gamma}(z_{a'}, Q^2) \frac{x_a x_b z_a z_{a'}}{x_a x_b z_{a'} - x_a z_{a'} x_2} \\ \times \frac{d\hat{\sigma}}{d\hat{t}}(x_a, x_b, z_a, z_{a'}, P_T), \end{aligned} \tag{12}$$

where $z_{a'}$ is the momentum fraction of partons from the resolved photon, and $F_{q_{a'}/\gamma}(z_{a'}, Q^2)$ is the parton distribution of the resolved photon [21]. The minimum values of momentum fractions are

$$x_{amin} = \frac{x_1}{1 - x_2}, \tag{13}$$



FIG. 1. The cross section of the real photon production in relativistic pp collisions at RHIC. The dash line is the sum of direct photons and fragmentation photons [15]. The dash dot line represents photons produced by the photoproduction and resolved photoproduction. The solid line is the sum of direct photons, fragmentation photons and photons from the photoproduction and resolved photoproduction. Data points are from PHENIX at RHIC [2].



FIG. 2. Same as Fig.1 but for pp $\sqrt{s}$=7 TeV collisions at LHC. Data points are from CMS at LHC [22].

and

$$x_{bmin} = \frac{x_a x_2}{x_a - x_1}, \tag{14}$$

and

$$z_{a'min} = \frac{x_b x_1}{x_a x_b - x_a x_2}. \tag{15}$$

The variable z_a of the resolved photon is given by

$$z_a = \frac{x_b x_1}{x_a x_b z_{a'} - x_a z_{a'} x_2}. \tag{16}$$

In the resolved photoproduction, the photon spectrum from the quark q_a can be written as

$$\begin{aligned} &f_{\gamma/q_a}(z_a) = \\ &\times \frac{e_q^2 \alpha P_{q\to\gamma}(z_a)}{2\pi}\left[\ln\left(\frac{(\sqrt{s_{NN}})^2}{4m_q^2}\right) + \ln\left(\frac{x_a x_b z_{a'}(1-z_a)}{z_a}\right)\right] \\ &- \frac{e_q^2\alpha}{\pi}\left(\frac{1-z_a}{z_a} - \frac{4m_q^2}{x_a x_b z_{a'} s_{NN}}\right). \end{aligned} \tag{17}$$

The cross section $d\hat{\sigma}/d\hat{t}$ of the annihilation and Compton scattering ($q_{a'}\bar{q}_b \to g\gamma$, $\bar{q}_{a'}q_b \to g\gamma$, $q_{a'}g_b \to q\gamma$, $g_{a'}q_b \to q\gamma$) are given by [1, 7]

$$\frac{d\hat{\sigma}}{d\hat{t}}(q\bar{q} \to g\gamma) = \frac{\pi\alpha\alpha_s e_q^2}{\hat{s}^2}\frac{8}{9}\left(\frac{\hat{u}}{\hat{t}} + \frac{\hat{t}}{\hat{u}}\right), \tag{18}$$

and

$$\frac{d\hat{\sigma}}{d\hat{t}}(qg \to q\gamma) = \frac{\pi\alpha\alpha_s e_q^2}{\hat{s}^2}\frac{1}{3}\left(-\frac{\hat{t}}{\hat{s}} - \frac{\hat{s}}{\hat{t}}\right), \tag{19}$$

where the strong coupling constant is

$$\alpha_s = \frac{12\pi}{(33-2n_f)\ln(Q^2/\Lambda^2)}, \tag{20}$$

the large momentum variable is $Q^2 = 4P_T^2$ and the momentum scale is Λ_{LO}= 0.2 GeV [7]. n_f is the flavor number. The Mandelstam variables in resolved photoproduction are

$$\hat{s} = x_a x_b z_a z_{a'} s_{NN}, \tag{21}$$

and

$$\hat{t} = -x_a x_2 z_a z_{a'} s_{NN}, \tag{22}$$

and

$$\hat{u} = -x_b x_1 s_{NN}. \tag{23}$$

In Fig.1 and 2 we plot the contribution of prompt photons and photons produced from photoproductions for relativistic pp collisions at RHIC and LHC energies. Except the leading QCD Compton and annihilation subprocesses, the pure QED annihilation $q\bar{q} \to \gamma\gamma$ (res.pho.) and QCD-induced gluon-photon coupling $g\gamma \to g\gamma$ (pho.), $g\gamma \to \gamma\gamma$ (pho.), $gg \to g\gamma$ (res.pho.) and $gg \to \gamma\gamma$ (res.pho.) [23] are considered in the subprocesses. The correction factors to take into account the NLO effect of subprocesses of direct photons and fragmentation photons are $K_{dir.\gamma}$ and $K_{fra.\gamma}$. We evaluate these P_T-dependent K factors using the numerical program from Aurenche et al [11]. The authors of Ref. [10] have obtained $K_{dir.\gamma}$(10 GeV)$\sim$ 1.5 for RHIC and LHC and $K_{fra.\gamma}$(10 GeV)$\sim$ 1.8 at RHIC and 1.4 at LHC.

In Fig.1 the spectra of the photoproduction and resolved photoproduction (dash dot line) are compared with the spectra of direct photons and fragmentation photons (dash line). The numerical results show that the modification of photoproductions is not prominent in the relatively small P_T region. The modification of the photoproductions is $\left(Ed\sigma_{pho.+res.pho.}/d^3P\right) / \left(Ed\sigma_{dir.+fra.}/d^3P\right) \sim 12\%$ in the region of 5 GeV $< P_T <$ 16 GeV at RHIC energies. However, the photoproductions start playing an interesting role in large P_T region at LHC energies. The modification in the region of 20 GeV $< P_T <$ 200 GeV is almost 33% at LHC (Fig.2).

In conclusion, the photoproductions are very important in the ep deep inelastic collisions at HERA, we extend the photoproduction mechanism to the real photon production in the pp collisionswe at RHIC and LHC energies. Based on the Weizsäcker-Williams approximation, we derive the high energy photon spectrum from the charged partons, and study the cross section of the photoproduction and resolved photoproduction. The photon spectrum from the charged parton depends on the collision energies, and the photoproductions can produce large P_T photons at RHIC and LHC. The numerical results show that the photoproductions contribute an active modification for the direct photons and fragmentation photons production. The modification of photoproductions is weak in the relatively small P_T, but becomes important in the large P_T region.

This work is supported by the National Natural Science Foundation of China (10665003 and 11065010).

---

[1] Owens J F 1987 *Rev. Mod. Phys.* **59** 465

[2] Peressounko D, the PHENIX Collaboration 2007 *Nucl. Phys.* A **783** 577

[3] Fu Y P and Li Y D 2011 *Nucl. Phys.* A **865** 76

[4] Fu Y P and Li Y D 2010 *Chin. Phys. Lett.* **27** 101202

[5] Fu Y P and Li Y D 2009 *Chin. Phys. Lett.* **26** 111201

[6] Fu Y P and Li Y D 2010 *Chin. Phys.* C **34** 186

[7] Field R D 1989 *Applications of Perturbative QCD* (New York: Addison-Wesley Publishing Company) pp 186-195

[8] Kang Z, Oiu J and Vogelsang W 2009 *Phys. Rev.* D **79** 054007

[9] Peitzmann T and Thoma M H 2002 *Phys. Rep.* **364** 175

[10] Turbide S, Gale C, Jeon S and Moore G D 2005 *Phys. Rev.* C **72** 014906

[11] Aurenche P, Baier R, Douiri A, Fontannaz M and Schiff D 1987 *Nucl. Phys.* B **286** 553

[12] Nisius R 2000 *Phys. Rep.* **332** 165

[13] Krawczyk M,Zembrzuski A and Staszel M 2001 *Phys. Rep.* **345** 265

[14] Forshaw J R arXiv:hep-ph/9706319

[15] Fu Y P and Li Y D 2011 *Phys. Rev.* C **84** 044906

[16] Yin R T, Fu Y P and Li Y D 2011 *Cent. Eur. J. Phys.* **9** 1434

[17] Baur G 1998 *J. Phys.* G **24** 1657

[18] Eboli O J P, Gonzalez-Garcia M C and Novaes S F 1994 *Phys. Rev.* D **49** 91

[19] Drees M, Godbole R M, Nowakowski M and Rindani S D 1994 *Phys. Rev.* D **50** 2335

[20] Glück M, Reya E and Vogt A 1992 *Z. Phys.* C **53** 127

[21] Glück M, Reya E and Vogt A 1992 *Phys. Rev.* D **46** 1973

[22] Khachatryan V, the CMS Collaboration 2011 *Phys. Rev. Lett.* **106** 082001

[23] Berger E L, Braaten E and Field R D 1984 *Nucl. Phys.* B **239** 52